

Early Control of Tough Broadleaf, Aquatic and Sedge Weeds

League[®] Herbicide is an unprecedented early season, lead-off play against tough broadleaf, aquatic and sedge weeds. It offers solid preemergence residual control of yield-robbing weeds and will fit your herbicide program lineup.

- ▶ **Strikes weeds as they emerge**, providing unique residual control for use on conventional or Clearfield[®] rice
- ▶ **Controls tough rice weeds**, including jointvetches, hemp sesbania, Texasweed, morningglory, flat sedge, eclipta, yellow nutsedge and more

How To Use

Preemergence Use Recommendations	<ul style="list-style-type: none"> • Rate: <i>League</i> 6.4 oz/A plus foundation grass herbicide (such as Bolero[®] 8 EC Rice Herbicide, Command[®] or Newpath[®]) • Residual broadleaf and sedge control—yellow nutsedge, early indigo (jointvetch), coffeebean and Texasweed control • Good approach for broadleaf control around towns and soybeans • Not for use on clay soils with pH >8.0 • Apply <i>League</i> to a well-prepared, moist seedbed. Soil should be sealed by flushing or rainfall prior to application of <i>League</i>.
Method	Ground or air
Spray Volume	Minimum 10 gal/A
REI	12 hours

Proven Control of Tough Weeds

League controls tough weeds and is a flexible fit in your weed management program.

Application rates: *League* 6.4 oz, Command 12.8 oz, Roundup[®] 32 oz and Grounded[®] 12.8 oz. Photos were taken 40 days after application.

Location: Winnie, TX.
Source: Valent



Command + Roundup + Grounded



League + Command + Roundup + Grounded

Weeds Controlled By *League* Plus Foundation Grass Herbicide

Broadleaves & Aquatics			
Croton, Woolly (<i>Croton capitatus</i>)	Gourd, Texas (<i>Cucurbita foetidissima</i>)	Jointvetch, Indian (<i>Aeschynomene indica</i>)	Pigweeds (except Livid) (<i>Amaranthus</i> spp.)
Ducksalad (<i>Heteranthera</i> spp.)	Groundcherry, cutleaf (<i>Physalis angulata</i>)	Jointvetch, Northern (<i>Aeschynomene virginica</i>)	Redstem (<i>Ammannia</i> spp.)
Eclipta (<i>Eclipta prostrata</i>)	Hemp Sesbania (<i>Sesbania exaltata</i>)	Morningglory, pitted (<i>Ipomoea lacunosa</i>)	Texasweed (<i>Caperonia palustris</i>)
Grasses			
Tank mix with a foundation grass herbicide, such as <i>Bolero</i> , <i>Command</i> , <i>Newpath</i>			
Barnyardgrass (<i>Echinochloa crus-galli</i>)	Crabgrass, large (<i>Digitaria sanguinalis</i>)	Panicum, Fall (<i>Panicum dichotomiflorum</i>)	Sprangletop (<i>Leptochloa</i> spp.)
Broadleaf signalgrass (<i>Brachiaria platyphylla</i>)	Johnsongrass (seedling) (<i>Sorghum halepense</i>)	Panicum, Texas (<i>Panicum texanum</i>)	
Sedges			
Dayflower (<i>Commelina communis</i>)	Flatsedge, rice (<i>Cyperus iria</i>)	Nutsedge, yellow (<i>Cyperus esculentus</i>)	

Other Important Information

- ▶ *League* is approved for aerial application. Do not apply *League* by air within 0.5 miles of emerged cotton or non-STS soybeans.
- ▶ *League* can be tank mixed with many rice herbicides, except Ricestar® HT and Clincher®
- ▶ Fields with a history of weed resistance to Londax® may exhibit resistance to *League*
- ▶ Do not apply more than 6.4 oz of *League* per acre per year