

We've Got Your Fields Covered, So Weeds Won't.

Bolero® 8 EC Herbicide applied preplant, late preemergence or postemergence controls many yield-impacting weeds in rice, including herbicide-resistant barnyardgrass. *Bolero* 8 EC can also provide residual control of some weeds for up to 5 weeks following application.

- ▶ Controls tough-to-manage barnyardgrass, plus aquatics and annual sedges
- ▶ Ideal in conventional or Clearfield® rice production
- ▶ Great fit in dry-seeded or water-seeded rice production
- ▶ Compatible with other delayed preemergence applications



How To Use

Rate	4 pt/A
Timing	<p>Water-Seeded—Red Rice Suppression and Sprangletop Control (Preplant, Non-Incorporated)</p> <ul style="list-style-type: none"> • Apply to a well-prepared seedbed • Flood field 2–3 days after application <p>Drill-Seeded (Delayed Preemergence)</p> <ul style="list-style-type: none"> • Apply to a well-prepared, moist seedbed • Rice seed must germinate (have a primary root at least ½ inch long) prior to application <p>Dry-Seeded (Postemergence)</p> <ul style="list-style-type: none"> • Apply to moist soil or flooded fields • Postemergence application to drill seeded rice can be made after emergence <p>Water-Seeded (Postemergence)</p> <ul style="list-style-type: none"> • Apply to moist soil or flooded fields • Postemergence application may be made to rice that is in at least the 2-leaf (second leaf fully expanded) stage of growth
Method	Air or ground
Spray Volume	<p>Air: Apply in no less than 10 gallons spray mixture per acre</p> <p>Ground: Apply in a minimum of 10 gallons of total spray mixture per acre</p>
REI	12 hours



Other Important Information

- ▶ Do not apply to fields with exposed seed as exposed seed will be killed
- ▶ Do not apply to stressed rice
- ▶ Do not apply as a preemergence treatment to cracked soil
- ▶ Do not apply more than 2 ½ pint per acre when using aerial application equipment east of the Rocky Mountains
- ▶ Do not apply more than 4 lb active ingredient per acre per year
- ▶ Do not apply more than 1 application per acre per year
- ▶ Do not apply through any type of irrigation system
- ▶ Do not release permanent flood water within 14 days of application of this product (where weather permits) when applying to rice fields
- ▶ Do not apply this product within 24 hours of rainfall, or when heavy rain is expected to occur within 24 hours
- ▶ Do not apply when temperatures exceed 95°F
- ▶ Do not overlap or double spray ends of field
- ▶ Do not apply to a second stubble rice crop
- ▶ Do not apply in combination with propanil within 14 days before or after organophosphate or carbamate insecticide application

