

## Add Speed and Efficiency to Your Tank with Resource® Herbicide

As the number of tough-to-control weeds increases, the demand for an effective tank mix partner also increases. Resource® Herbicide is a reliable choice as a tank mix partner with glyphosate. *Resource* is a powerful and cost-effective solution for growers with problem weeds like velvetleaf, morningglory, common ragweed, pigweeds, smartweeds and lambsquarters.

- ▶ Postemergence herbicide partner
- ▶ Kills problem weeds that glyphosate alone misses
- ▶ Adds speed to glyphosate

### How To Use—Field Corn\*

<b>Application Rate</b>	<b>Broadcast:</b> 4–6 fl oz/A <b>Drop-Nozzle:</b> 4–8 fl oz/A <b>Tank Mix:</b> 4–8 fl oz/A <b>Tank Mix with glyphosate for increased control of velvetleaf:</b> 2–4 fl oz/A
<b>Restrictions</b>	<ul style="list-style-type: none"> <li>• Do not apply more than 6 fl oz/A in a single broadcast application or more than a total of 8 fl oz/A to field corn during a single growing season</li> <li>• As a directed spray using drop nozzles, do not apply more than a total of 8 fl oz /A to field corn in a single application or during a single growing season</li> </ul>
<b>Application Timing</b>	May be applied to field corn from the 2-leaf through the 10-leaf stage
<b>Spray Volume</b>	Minimum of 20 gal/A
<b>Adjuvant</b>	<ul style="list-style-type: none"> <li>• Must be applied with 1 pt/A of crop oil concentrate or methylated seed oil. A spray grade nitrogen source (either ammonium sulfate at 2–2.5 lb/A or 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to enhance weed control.</li> <li>• For tank mixes, refer to tank mix partner’s label for adjuvant recommendation</li> </ul>
<b>Rotation Restrictions</b>	Do not rotate to crops other than field corn or soybeans within 30 days after last application
<b>Hay / Grazing Restrictions</b>	Do not graze animals on green forage or use as feed fewer than 28 days after application
<b>REI</b>	12 hours

\*Do not use *Resource* on popcorn or sweet corn.

## How To Use—Soybeans\*\*

<b>Application Rate</b>	<b>Broadcast:</b> 4–8 fl oz/A (8–12 fl oz/A for late control of tall velvetleaf) <b>Tank Mix:</b> 4–12 fl oz/A <b>Tank Mix with glyphosate for increased control of velvetleaf and suppression of morningglories:</b> 2–4 fl oz/A
<b>Restrictions</b>	<ul style="list-style-type: none"> <li>Do not apply more than 12 fl oz /A in a single application or more than a total of 16 fl oz /A to soybeans during a single growing season</li> <li>Do not apply if rain is expected within one hour of application; otherwise unsatisfactory weed control may result</li> </ul>
<b>Application Timing</b>	Burndown or postemergence up to 60 days prior to harvest (follow restrictions associated with tank mix partner)
<b>Spray Volume</b>	Minimum of 20 gal/A
<b>Adjuvant</b>	<ul style="list-style-type: none"> <li>Must be applied with 1 qt/A of crop oil concentrate or methylated seed oil. A spray grade nitrogen source (either ammonium sulfate at 2–2.5 lb/A or 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to enhance weed control.</li> <li>For tank mixes, refer to tank mix partner’s label for adjuvant recommendation</li> </ul>
<b>Rotation Restrictions</b>	Do not rotate to crops other than field corn or soybeans within 30 days after last application
<b>Hay / Grazing Restrictions</b>	Do not graze treated fields or harvest for forage or hay
<b>REI</b>	12 hours
<b>PHI</b>	60 days

\*\*Resource is not for use in California on soybeans.

