


Quash® Fungicide Controls White Mold, Boosts Yield Potential

Quash® Fungicide is a highly effective foundation fungicide that delivers proven, broad spectrum performance for a planned rotation program. *Quash* controls white mold, maximizing canola yield potential with an excellent return on investment (ROI).

- ▶ Controls White Mold/Sclerotinia Stem Rot (*Sclerotinia sclerotiorum*)
- ▶ Maximizes Yield Potential
- ▶ Proven Results—Improved Quality and ROI

Quash Controls Severity of White Mold in Canola



How To Use

| | |
|---------------------|---|
| Rate | <ul style="list-style-type: none"> • 2–4 oz/A (use 4 oz/A under high disease pressure) • Section 2(ee) for Minnesota, Montana (east of Interstate 15 only) and North Dakota: 1.5–4 oz/A + 10–20 fl oz/A fungicide containing thiophanate-methyl as a single active ingredient and labeled for use in canola |
| Timing | Apply between 20% and 50% bloom |
| Method | Ground, aerial, chemigation |
| Spray Volume | 10–20 gal/A for ground application and 5 gal/A for aerial application |
| Adjuvant | Not required, but allowed |
| Restrictions | <ul style="list-style-type: none"> • Do not apply more than 4 oz of product per acre per year • Do not make more than one application per year |
| REI / PHI | 35 days / 12 hours |

